

Spatiotemporal decoupling of littoral and lacustrine geosmin dynamics: Implications for early warning in drinking water reservoirs

Highlights

Yuying Gui^{a,b,#} Tengxin Cao^{b,f,#} Jie Yang^c Jihui Qin^d Ming Su^{*,b,f,*} Ziyi Yang^{b,f}
Qi Zhang^e Yufan Ai^{b,f} Jiao Fang^b Yingjie Li^{b,f} Yuanhong Xiao^d Zhixiang Hao^c
 Zhengyan Li^a Min Yang^{b,f}

These authors contributed equally to this work.

^a College of Environmental Science and Engineering, Ocean University of China, Qingdao 266100, China.

^b Key Laboratory of Environmental Aquatic Chemistry, State Key Laboratory of Regional Environment and Sustainability, Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences, Beijing 100085, China.

^c Tianjin Hydraulic Research Institute, Tianjin 300061, China.

^d Tianjin Yuqiao Reservoir Authority, Tianjin 301900, China.

^e Institute of Hydrobiology, Chinese Academy of Sciences, Wuhan 430070, China.

^f University of Chinese Academy of Sciences, Beijing 100049, China.

* Corresponding to: [Ming Su \(mingsu@rcees.ac.cn\)](mailto:mingsu@rcees.ac.cn)

- Unveils littoral zone as the critical initiation hotspot for reservoir odor
- Develops a dynamic early-warning model integrating multi-source sensing data
- Proposes targeted intervention protocol based on odor source-pathway identification
- Validates framework via cross-seasonal, multi-reservoir field campaigns