

MIB-derived odor management based upon hydraulic regulation in small drinking water reservoirs: principle and application

Jinping Lu^{a,d}, Ming Su^{a,d,*}, Yuliang Su^b, Jiao Fang^a, Michael Burch^c, Tengxin Cao^{a,d}, Bin Wu^b, Jianwei Yu^{a,d}, Min Yang^{a,d,*}

^aState Key Laboratory of Environmental Aquatic Chemistry, Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences, P.O. Box 2871, Beijing, 100085,

^bZhuhai Water Environment Holdings Group Ltd., Zhuhai, 519020,

^cSchool of Biological Sciences, The University of Adelaide, SA, 5005,

^dUniversity of Chinese Academy of Sciences, Beijing, 100049,

Highlights

- Hydraulic retention time (HRT) increase can trigger 2-methylisoborneol (MIB) episodes
- MIB-producing *Planktothricoides* can be restricted by reducing HRT to <10 days
- Critical HRTs were evaluated for 14 cyanobacterial genera based on a theoretical model
- HRT regulation was successfully applied for MIB control in ZXD Reservoir

*Corresponding author

Email addresses: mingsu@rcees.ac.cn (Ming Su), yangmin@rcees.ac.cn (Min Yang)