

# Biosynthesis of 2-methylisoborneol is regulated by chromatic acclimation of *Pseudanabaena*

Ming Su<sup>a,b,g,\*</sup>, Jiao Fang<sup>a,h</sup>, Zeyu Jia<sup>a,c</sup>, Yuliang Su<sup>d</sup>, Yiping Zhu<sup>e</sup>, Bin Wu<sup>d</sup>, John C. Little<sup>f</sup>, Jianwei Yu<sup>a,b,g</sup>, Min Yang<sup>a,b,g</sup>

<sup>a</sup>Key Laboratory of Drinking Water Science and Technology, Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences, P.O. Box 2871, Beijing, 100085,

<sup>b</sup>National Engineering Research Center of Industrial Wastewater Detoxication and Resource Recovery, Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences, P.O. Box 2871, Beijing, 100085,

<sup>c</sup>Yangtze Eco-Environment Engineering Research Center, China Three Gorges Corporation, Beijing, 100038,

<sup>d</sup>Zhuhai Water Environment Holdings Group Ltd., Zhuhai, 519020,

<sup>e</sup>Shanghai Chengtou Raw Water Co. Ltd., Beiai Rd. 1540, Shanghai, 200125,

<sup>f</sup>Department of Civil and Environmental Engineering, Virginia Tech., Blacksburg, VA, 24061-0246,

<sup>g</sup>University of Chinese Academy of Sciences, Beijing, 100049,

<sup>h</sup>School of Civil Engineering, Chang'an University, Xi'an, 710054,

---

## Highlights

- Biosynthesis of 2-methylisoborneol (MIB) is regulated by chromatic acclimation
- MIB is correlated with chlorophyll *a* abundance regardless with light color
- Green light promote MIB synthesis of phycoerythrin-containing *Pseudanabaena*

## References

---

\*Corresponding author  
Email addresses: mingsu@rcees.ac.cn (Ming Su), jia\_zeyu@ctg.com.cn (Zeyu Jia), jwyu@rcees.ac.cn (Jianwei Yu), yangmin@rcees.ac.cn (Min Yang)